

MATHEMATICS

MATH-101-E

Time : Three Hours]

[Maximum Marks : 100

Note : Attempt *Five* questions in all, selecting at least *one* question from each Unit. All questions carry equal marks.

Unit I

1. (a) Compute to four decimal places, the value of $\cos 32^\circ$ by use of Taylor's series.

(b) Define asymptotes of a polar curve. Find asymptotes of the curve $r \sin \theta = 2 \cos 2\theta$.

2. (a) Find the radius of curvature for the parabola

$$\frac{2a}{r} = 1 + \cos \theta.$$

(b) Trace the curve :

$$y^2 (a - x) = x^2 (a + x)$$

Unit II

23 (a) Define Homogeneous function of degree n . State and prove Euler's Theorem for a homogeneous function of degree n in x , y and z .

(b) If $x = e^u \cos v$, $y = e^u \sin v$, find

$$J = \frac{\partial(u, v)}{\partial(x, y)}, \quad J' = \frac{\partial(x, y)}{\partial(u, v)} \text{ and hence show}$$

that :

$$JJ' = 1.$$

4. (a) Show that the rectangular solid of maximum volume that can be inscribed in a sphere is a cube.

(b) Evaluate $\int_0^\pi \log(1 + a \cos x) dx$ using the method of differentiation under the sign of integration.

Unit III

5. (a) Find by double integration, the area lying inside the circle $r = a \sin \theta$ and outside the cardioids $r = a(1 - \cos \theta)$.

- (b) Evaluate $\int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy$ by changing into polar co-ordinates. Hence show that :

$$\int_0^{\infty} e^{-x^2} dx = \sqrt{\frac{\pi}{2}}$$

6. (a) Find the volume of the tetrahedron bounded by the co-ordinate planes and the plane

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1.$$

- (b) Define Beta and Gamma functions. Prove that :

$$\beta\left(m, \frac{1}{2}\right) = 2^{2m-1} \beta(m, m)$$

Unit IV

7. (a) Define Gradient of a scalar point function and hence give its geometrical interpretation.

(15)

- (b) With usual notations, prove that :

(15)

$$\nabla^2 (r^n) = n(n+1)r^{n-2}$$

8. (a) Verify Green Theorem for

$$\int_C \left[(3x - 8y^2) dx + (4y - 6xy) dy \right] \text{ over } C,$$

the boundary of the region bounded by $x = 0$, $y = 0$ and $x + y = 1$.

(b) Evaluate :

$$\int \bar{F} \cdot d\bar{S}$$

over the surface S bounding by the region

$x^2 + y^2 = 4$, $z = 0$ and $z = 3$ by taking

$$\bar{F} = 4xi - 2y^2j + z^2k.$$